

Reference values of aortic pulse wave velocity in a large healthy population aged between 3 and 18 years

Erzsébet Valéria Hidvégi^a, Miklós Illyés^a, Béla Benczúr^a, Renáta M. Böcskei^a, László Rátgéber^b, Zsófia Lenkey^a, Ferenc T. Molnár^c, and Attila Cziráki^a

Objective: The measurement of aortic pulse wave velocity (PWV_{ao}) is an accepted marker in stratifying individual cardiovascular risk in adults. There is an increasing volume of evidence concerning impaired vascular function in different diseases in paediatric populations, but, unfortunately, only a few studies are available on the measurement of normal PWV_{ao} values in children. The aim of our study was to determine the reference values of PWV_{ao} in a large healthy population using a newly developed technique.

Methods: Three thousand, three hundred and seventy-four healthy individuals (1802 boys) aged 3–18 years were examined by an invasively validated, occlusive, oscillometric device.

Results: The mean PWV_{ao} values increased from 5.5 ± 0.3 to 6.5 ± 0.3 m/s ($P < 0.05$) in boys and from 5.6 ± 0.3 to 6.4 ± 0.3 m/s ($P < 0.05$) in girls. The increase, however, was not constant, and the values exhibited a flat period between the ages of 3 and 8 years in both sexes. The first pronounced increase occurred at the age of 12.1 years in boys and 10.4 years in girls. Moreover, between the ages of 3 and 8 years, the brachial SBP and mean blood pressures increased continuously and gradually, whereas the PWV_{ao} remained unchanged. By contrast, beyond the age of 9 years, blood pressure and aortic stiffness trends basically moved together.

Conclusion: Our study provides the largest database to date concerning arterial stiffness in healthy children and adolescents between the ages of 3 and 18 years, and the technology adopted proved easy to use in large paediatric populations, even at a very young age.

Keywords: aorta, Arteriograph, blood pressure, children and adolescents, pulse wave velocity

Abbreviations: CCA, common carotid artery; HR, heart rate; MAP, mean arterial pressure; PWV_{ao}, aortic pulse wave velocity; RCA, right carotid artery; SSN, suprasternal notch; TEM, technical errors of measurement

individual cardiovascular risk in adults [1]. Moreover, we have an increasing volume of evidence concerning impaired vascular function in different diseases in paediatric populations: early atherosclerosis [2], obesity [3], extreme prematurity [4], familial hypercholesterolemia [5,6], type I diabetes mellitus [7,8], juvenile hypertension [9], different congenital heart diseases [10,11], end-stage renal disease [12], HIV infection [13], Kawasaki disease [14], neurofibromatosis [15], vasculitis [16], intrauterine growth restriction patients [17] and primary snoring [18].

These diseases might well influence aortic stiffness, and so the measurement of PWV_{ao} may provide useful information, even in a paediatric population. However, to be able to assess the measured PWV_{ao} values in different conditions and, further, to assess age-related dynamic changes of PWV_{ao} in children and adolescents – especially in younger populations of less than 6 years – a precise determination of reference values between the third and 97th percentile is extremely important.

Unfortunately, few studies are available which detail the normal values of PWV_{ao} in paediatric populations. To date, in fact, a total of 1514 healthy individuals have been studied in six articles [19–24] (Table 1). In the majority of these studies, relatively small populations ($n < 140$) were measured to provide control groups to the diseased populations studied, and it is also worth mentioning that these healthy control groups were remarkably imbalanced in terms of age distribution. Only one study [24] comprised a larger population (1008 individuals, 6–20 years), but even in this study the age distribution of the population was poorly balanced, as a substantial majority (68.1%) of the individuals studied were from the 15–20 years age group. On the contrary, these studies used ultrasound and applanation tonometry to determine PWV_{ao}, techniques which are limited in use

Journal of Hypertension 2012, 30:2314–2321

^aHeart Institute, Faculty of Medicine, University of Pécs, ^bFaculty of Health Sciences, Institute of Public Health and Health Promotion, University of Pécs, Pécs and ^cDepartment of Hydrodynamic Systems, Budapest University of Technology and Economics, Budapest, Hungary

Correspondence to Erzsébet Valéria Hidvégi, MD, Heart Institute, Faculty of Medicine, University of Pécs, 13 Ifjúság str., H-7624 Pécs, Hungary. Tel/fax: +36 56 522 301; e-mail: pediatriccor@gmail.com

Received 15 January 2012 **Revised** 24 May 2012 **Accepted** 14 August 2012

J Hypertens 30:2314–2321 © 2012 Wolters Kluwer Health | Lippincott Williams & Wilkins.

DOI: 10.1097/HJH.0b013e328359562c

INTRODUCTION

The measurement of arterial stiffness, that is, aortic pulse wave velocity (PWV_{ao}), is an accepted marker for detecting organ damage and for stratifying

TABLE 1. Published normal paediatric aorta pulse wave velocity values

Age, years	Number of participants	Method	Distance measurement	PWV _{ao} , m/s	Reference
11	30	Pressure transducers	CCA—a. fem.	6.5 ± 1.2	Tedesco <i>et al.</i> [19]
12.9 ± 0.2	110	Applanation tonometry	RCA—right a. fem	4.1–8.2	Ahimastos <i>et al.</i> [20]
0.2–20	125	Doppler ultrasound	CCA—a. fem.	4.4–7.9	Avolio <i>et al.</i> [21]
3–20	108	Doppler ultrasound	CCA—a. fem.; MS—a. fem	6.2–10.2	Avolio <i>et al.</i> [22]
6–23	133	Applanation tonometry	CCA—SSN+SSN—a. fem	5.02 ± 0.89	Kis <i>et al.</i> [23]
6–20	495 male; 513 female	Applanation tonometry	CCA—SSN –SSN—a. fem	Male 4.34–5.71; female 4.34–5.50	Reusz <i>et al.</i> [24]

a.fem, femoral artery; CCA, common carotid artery; MS, manubrium sterni; RCA, right carotid artery; SSN, suprasternal notch.

to the noninvasive clinical evaluation of aortic stiffness in paediatric patients [25].

The aim of our study, therefore, was to determine the reference values of the PWV_{ao} in a large healthy population which showed a fair balance in terms of age. We intended to apply a newly developed, invasively validated and easy-to-use oscillometric technique which allowed us to expand measurement even into a very young (<6 years) population and within a daily clinical routine.

METHODS

Participants

Three thousand, three hundred and seventy-four apparently healthy white individuals without any medication and/or blood pressure-affecting drugs from Hungary (1802 boys and 1572 girls) aged 3–18 years, having BMI and SBP and DBP within the third and 97th percentiles, as established by the relevant guidelines [26,27], were recruited from nursery, elementary and secondary schools. Written consent for the measurements to be taken was given by the parents. The protocol of this clinical study was reviewed and approved by the local Institutional Ethics Committee of the University of Pécs, Pécs, Hungary.

Methods

A noninvasive, occlusive, oscillometric device (Arteriograph; TensioMed Ltd, Budapest, Hungary) was used for the PWV_{ao} measurements, for which the operating procedure did not, in practice, differ from a standard digital blood pressure measurement. The method and its validations have been detailed previously [28,29]. Briefly, the method is based on the physiological fact that the early (P₁) systolic pulse pressure wave of the aorta, generated by the left ventricle ejection, travels along the aorta and is reflected (P₂) from the area of the aortic bifurcation. Occluding the brachial artery by pressurizing the cuff 35–40 mmHg above the actual SBP creates easily distinguished, pronounced

pressure peaks in the cuff. Separated in this way, early and late systolic waves can be recorded. The time lapses between the peaks of P₁ and P₂ are equal to the travel time of the aortic pressure wave from the aortic root to the bifurcation and back. By halving this time and measuring the sternal notch – pubic bone distance (which is rather close to the true aortic length [30]), the PWV_{ao} can be calculated (PWV_{ao} = jugulum–symphysis distance/transit time). The PWV_{ao} measurements were taken with the participant in a calm, supine position after several minutes of rest.

The interobserver and intraobserver errors of the distance (sternal notch–pubic bone) measurements were expressed as absolute and relative technical errors of measurement (TEM) [31] and proved to be well within acceptable limits (Table 2).

Data analysis and statistics

Data are reported as mean and SD for continuous data. For data comparison, a Student's *t*-test was used after checking that the assumption of normality was met. A significance level of 0.05 was used for statistical tests. Statistical analysis was performed with the SPSS 15.0 statistical package (SPSS Inc., Chicago, Illinois, USA).

Reference percentile curves were obtained with the use of the LMS method [32], specifically with the LMS Chartmaker software [33]. The method summarizes the distribution of a variable by its skewness (expressed as Box–Cox power, L), its median (M) and its coefficient of variation (S). The age-dependent changes in these parameters are smoothed by fitting cubic splines with appropriately selected smoothing parameters, using penalized likelihood.

RESULTS

The characteristics of the subjects investigated are summarized in Tables 3 and 4. Both the boys and the girls are fairly

TABLE 2. Intraobserver and interobserver errors

Physician(s)	Intraobserver					Interobserver									
	1	2	3	4	5	1–2	1–3	1–4	1–5	2–3	2–4	2–5	3–4	3–5	4–5
TEM (cm)	0.23	0.31	0.26	0.28	0.20	0.27	0.27	0.34	0.30	0.30	0.32	0.32	0.32	0.27	0.30
Relative TEM (%)	0.5	0.7	0.5	0.6	0.4	0.6	0.6	0.7	0.6	0.6	0.7	0.7	0.7	0.6	0.6

TEM, technical error of measurement.

TABLE 3. Characteristics of boys

Age (years)	n	Height (cm)	Weight (kg)	BSA (m ²)	BMI (%)	SBP _{brach} (mmHg)	DBP _{brach} (mmHg)	MAP (mmHg)	HR (beats/min)	JUG-SY (cm)
03	044	104.4±6.7	16.2±02.5	0.7±0.1	14.8±1.9	103.0±5.5	61.8±7.3	76.5±7.5	98.7±11.1	32.0±2.5
04	053	108.6±7.6	17.5±02.7	0.7±0.1	14.8±1.6	104.1±7.7	61.5±7.5	76.9±8.5	92.5±13.0	33.1±2.9
05	080	115.5±6.8	19.7±02.9	0.8±0.1	14.8±1.7	106.4±7.2	62.7±6.3	78.3±7.8	91.5±13.4	35.0±2.2
06	120	123.2±6.2	22.4±03.5	0.9±0.1	14.7±1.6	104.5±6.8	62.0±6.3	76.3±6.4	86.6±11.5	37.2±2.7
07	085	127.8±6.8	25.2±04.2	0.9±0.1	15.4±1.8	107.8±7.2	63.8±6.3	78.8±6.7	83.7±13.1	39.0±2.7
08	074	132.9±8.0	27.9±04.7	1.0±0.1	15.8±1.8	106.9±7.1	62.8±5.3	77.4±5.4	79.9±13.2	40.0±2.9
09	092	138.5±7.4	31.2±05.9	1.1±0.1	16.1±1.9	109.3±6.9	64.8±6.8	79.2±6.1	79.7±13.5	42.0±3.3
10	081	144.5±6.6	36.1±06.5	1.2±0.1	17.2±2.2	111.2±7.3	64.7±5.5	80.5±5.7	78.7±12.3	44.0±3.4
11	080	148.4±8.2	38.9±07.8	1.3±0.2	17.5±2.4	111.5±7.5	65.2±5.7	80.7±5.9	73.7±11.5	44.3±3.3
12	101	155.3±9.6	44.3±09.3	1.4±0.2	18.2±2.5	115.0±7.5	65.3±6.3	82.1±6.1	77.5±13.8	47.2±4.2
13	169	163.1±9.5	50.8±09.3	1.5±0.2	19.0±2.3	117.2±7.8	64.8±6.3	82.3±6.1	76.3±12.6	49.2±3.7
14	187	169.0±9.8	55.5±09.8	1.6±0.2	19.3±2.4	119.5±7.4	65.7±5.4	83.7±5.2	76.2±13.2	51.0±3.4
15	171	174.1±7.6	59.9±09.2	1.7±0.2	19.7±2.4	121.1±8.0	67.5±6.2	85.4±6.0	71.3±12.8	51.8±3.4
16	162	175.9±7.1	64.3±11.2	1.8±0.2	20.7±2.8	124.5±8.5	68.6±6.2	87.3±6.2	72.5±14.3	53.1±3.3
17	197	178.0±7.2	67.5±10.2	1.8±0.2	21.3±2.7	126.3±8.6	68.8±6.5	88.0±6.2	69.9±11.9	54.5±3.4
18	106	179.6±6.9	69.6±08.7	1.9±0.1	21.6±2.4	128.2±8.3	67.4±7.3	87.7±6.4	70.3±13.2	55.3±3.5

Data are shown as mean ± SD. DBP_{brach}, brachial DBP; HR, heart rate; JUG-SY, distance between sternal notch (jugulum) and upper part of the pubic bone (symphysis); MAP, mean arterial pressure; SBP_{brach}, brachial SBP.

well balanced in terms of age, whereas 15% of the population was recruited from very young age groups (3–6 years).

All of the parameters measured showed a marked increase with age, except for the heart rate – which decreased, as physiologically expected.

Table 5 summarizes the PWV_{ao} changes. The increase of mean PWV_{ao} values turned out to be roughly 1 m/s in both sexes from the age of 3 to 18 years, as this parameter increased from 5.5 ± 0.3 to 6.5 ± 0.3 m/s ($P < 0.05$) in boys and from 5.6 ± 0.3 to 6.4 ± 0.3 m/s ($P < 0.05$) in girls. When comparing the mean PWV_{ao} values in each year between boys and girls, noticeable differences were found in the 14-year and 16-year age groups. However, these differences are merely 0.1 and 0.2 m/s, respectively, and so cannot be considered as clinically significant.

Smoothed percentile curves of PWV_{ao} from the third to the 97th are shown for boys and girls in Fig. 1, and the

corresponding values of PWV_{ao} can be seen in Tables 6 and 7.

Analysing the curves of the median values of PWV_{ao} in boys and girls, it became clear that the increase was not constant with age, but had more complex dynamics (Fig. 1). In both sexes, the PWV_{ao} values exhibit a flat period between 3 and 8 years, with a steeper increase thereafter. The first derivative of the smoothed median curve of PWV_{ao} showed that the first pronounced increase occurred at the age of 12.1 years in boys and 10.4 years in girls (Fig. 2).

To study causal relationships between PWV_{ao} and other variables, graphical analyses were used, as multivariate linear regression cannot be applied due to the high multicollinearity among age, anthropometric and hemodynamic parameters. In Fig. 3, we demonstrate the relationship between PWV_{ao}, mean arterial pressure (MAP) and brachial SBP (SBP_{brach}), featuring the median of the standardized

TABLE 4. Characteristics of girls

Age (years)	n	Height (cm)	Weight (kg)	BSA (m ²)	BMI (%)	SBP _{brach} (mmHg)	DBP _{brach} (mmHg)	MAP (mmHg)	HR (beats/min)	JUG-SY (cm)
03	035	102.2±06.3	15.5±2.4	0.7±0.1	14.8±1.2	102.5±5.1	61.1±9.4	75.4±8.1	101.1±14.4	30.9±1.9
04	043	107.1±06.4	17.1±2.9	0.7±0.1	14.8±1.8	103.3±5.2	60.0±5.2	75.2±4.9	092.4±11.6	31.8±2.8
05	044	114.5±07.4	18.8±3.3	0.8±0.1	14.3±1.7	103.7±6.3	61.7±7.2	76.1±6.9	093.7±11.7	33.4±2.1
06	068	123.0±06.6	22.2±4.0	0.9±0.1	14.6±1.7	105.0±6.9	62.1±5.8	76.8±6.7	087.3±13.3	35.9±2.4
07	072	128.6±10.3	25.2±7.1	0.9±0.2	15.0±2.3	106.4±7.3	63.2±5.6	77.7±6.0	086.5±12.5	38.2±3.3
08	039	133.9±07.1	28.1±5.7	1.0±0.1	15.6±2.4	107.6±7.3	63.6±4.6	78.3±5.0	083.2±11.1	39.2±2.3
09	064	138.6±08.1	31.9±6.5	1.1±0.1	16.5±2.3	109.6±6.4	64.8±5.8	79.3±5.4	084.9±11.7	40.8±3.1
10	063	144.0±07.1	34.4±6.6	1.2±0.1	16.5±2.3	111.1±7.3	64.8±5.5	80.2±5.5	082.5±09.4	42.4±3.2
11	049	150.8±09.8	39.5±8.4	1.3±0.2	17.2±2.5	112.1±6.8	65.4±5.0	81.0±5.0	080.1±09.4	44.2±3.7
12	085	156.8±07.8	44.6±8.8	1.4±0.2	18.0±2.5	113.4±7.5	65.6±5.4	81.5±5.6	079.5±12.8	46.2±3.5
13	195	160.8±07.7	49.6±7.9	1.5±0.1	19.1±2.6	115.1±7.6	65.4±6.2	82.0±5.9	078.6±13.8	47.8±3.0
14	189	163.7±06.5	52.3±8.3	1.5±0.1	19.5±2.6	117.6±8.2	67.0±6.2	83.8±6.3	077.7±14.2	48.7±2.8
15	181	164.8±08.0	54.8±9.3	1.6±0.2	20.1±2.7	117.2±7.6	68.5±5.8	84.7±5.9	077.1±12.3	48.6±3.4
16	174	164.9±06.7	55.8±8.5	1.6±0.1	20.5±2.7	117.8±7.4	69.2±5.4	85.4±5.5	074.4±11.7	49.1±3.2
17	175	165.9±06.3	56.2±9.3	1.6±0.1	20.4±2.9	117.7±8.0	68.0±6.0	84.5±6.1	074.7±10.2	49.8±3.1
18	096	166.3±06.6	57.5±8.4	1.6±0.1	20.8±2.8	118.8±8.6	68.2±6.0	85.1±6.3	077.0±11.0	51.2±2.8

Data are shown as mean ± SD. Data are shown as mean ± SD. DBP_{brach}, brachial DBP; HR, heart rate; JUG-SY, distance between sternal notch (jugulum) and upper part of the pubic bone (symphysis); MAP, mean arterial pressure; SBP_{brach}, brachial SBP.

TABLE 5. Normal aortic pulse wave velocity mean values in boys and girls

Age (years)	Boys		Girls		Student's <i>t</i> -test
	<i>n</i>	PWV _{ao} (m/s) ± SD	<i>n</i>	PWV _{ao} (m/s) ± SD	
03	044	5.5 ± 0.6	035	5.6 ± 0.6	NS
04	053	5.5 ± 0.6	043	5.3 ± 0.6	NS
05	080	5.5 ± 0.6	044	5.4 ± 0.8	NS
06	120	5.4 ± 0.7	068	5.3 ± 0.7	NS
07	085	5.5 ± 0.5	072	5.5 ± 0.6	NS
08	074	5.4 ± 0.6	039	5.4 ± 0.6	NS
09	092	5.6 ± 0.6	064	5.5 ± 0.6	NS
10	081	5.5 ± 0.7	063	5.7 ± 0.7	NS
11	080	5.6 ± 0.7	049	5.7 ± 0.7	NS
12	101	5.8 ± 0.7	085	5.8 ± 0.5	NS
13	169	5.8 ± 0.6	195	5.9 ± 0.6	NS
14	187	6.0 ± 0.7	189	5.9 ± 0.6	<i>P</i> < 0.05
15	171	6.0 ± 0.6	181	6.0 ± 0.6	NS
16	162	6.2 ± 0.6	174	6.0 ± 0.5	<i>P</i> < 0.004
17	197	6.3 ± 0.6	175	6.2 ± 0.5	NS
18	106	6.5 ± 0.5	096	6.4 ± 0.5	NS

PWV_{ao}, aortic pulse wave velocity.




FIGURE 1 Smoothed percentile curves from third to 97th of aortic pulse wave velocity (PWV_{ao}) related with age for boys (a) and girls (b).

values of the parameters mentioned in boys and girls. Interestingly, at ages between 3 and 8 years a marked difference was observed in the trend of the changes in SBP_{brach} and MAP versus PWV_{ao}. Although the blood pressure increased continuously, the PWV_{ao} remained practically unchanged in this age range. However, beyond 9 years of age, the blood pressure and aortic stiffness trends basically travelled together.

DISCUSSION

The most important finding of our study is that we were able to determine the reference values of aortic stiffness on the largest database to date of healthy children and adolescents between the ages of 3 and 18 years. Our homogeneous white population was fairly well balanced in terms of both age and sex.

More importantly, a special characteristic of our database is that it contains a large number of very young subjects (<6 years), as the applied oscillometric, occlusive method (Arteriograph) allowed us to perform the measurements even in this special population. The operational procedure of this device basically does not differ from a standard digital blood pressure measurement and lasts only 2 min. It is a very rapid and painless procedure and is well tolerated even by the youngest participants.

The small size of the portable device enabled us to visit kindergartens and schools to collect the database very easily. Consequently, the adoption of this technology for routine paediatric practice seems to be realistic, especially in view of the user independence of the method.

The measurement of PWV_{ao} of the used method is based on the assumption that the reflection site of the first systolic aortic pulse wave is at the area of the aortic bifurcation. Horvath *et al.* [29] clearly demonstrated that the PWV_{ao} measured invasively and oscillometrically were very close (<0.1 m/s) to each other, suggesting that the major reflection site must be at the area of the bifurcation. As far as the

TABLE 6. Aortic pulse wave velocity percentile data for boys

Age (years)	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
n	44	53	80	120	85	74	92	81	80	101	168	187	171	162	197	106
PWV _{ao} percentile data	L	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	M	5.477	5.478	5.479	5.482	5.483	5.492	5.514	5.549	5.683	5.796	5.911	6.010	6.116	6.240	6.382
	S	0.110	0.110	0.110	0.110	0.110	0.111	0.111	0.112	0.113	0.113	0.110	0.105	0.097	0.089	0.083
	97 th	6.69	6.69	6.69	6.69	6.69	6.71	6.74	6.86	6.97	7.11	7.22	7.28	7.31	7.35	7.44
	90 th	6.28	6.29	6.29	6.29	6.29	6.30	6.33	6.37	6.44	6.54	6.67	6.78	6.85	6.98	7.09
	75 th	5.88	5.88	5.88	5.89	5.89	5.90	5.92	6.02	6.11	6.23	6.35	6.43	6.51	6.61	6.73
	50 th	5.48	5.48	5.48	5.48	5.48	5.49	5.51	5.55	5.60	5.68	5.80	5.91	6.01	6.12	6.24
	25 th	5.07	5.08	5.08	5.08	5.08	5.09	5.11	5.18	5.25	5.36	5.48	5.59	5.72	5.87	6.03
	10 th	4.67	4.67	4.67	4.67	4.68	4.68	4.70	4.76	4.82	4.92	5.04	5.17	5.32	5.50	5.68
	3 rd	4.27	4.27	4.27	4.27	4.27	4.28	4.29	4.34	4.40	4.49	4.61	4.74	4.93	5.13	5.32
Height (cm)	104.4	108.6	115.5	123.2	127.8	132.9	138.5	144.5	148.4	155.3	163.1	169.0	174.1	175.9	178.1	179.6
Weight (kg)	16.2	17.5	19.7	22.4	25.2	27.9	31.2	36.1	38.9	44.3	50.8	55.5	59.9	64.3	67.6	69.6

PWV_{ao}, aortic pulse wave velocity.

TABLE 7. Aortic pulse wave velocity percentile data for girls

Age (years)	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
n	35	43	44	68	72	39	64	63	49	85	196	189	181	174	175	96
PWV _{ao} percentile data	L	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
	M	5.384	5.398	5.407	5.411	5.426	5.456	5.581	5.675	5.763	5.838	5.901	5.957	6.019	6.126	6.290
	S	0.123	0.122	0.122	0.122	0.121	0.120	0.118	0.113	0.109	0.106	0.102	0.097	0.091	0.084	0.077
	97 th	6.71	6.72	6.72	6.73	6.74	6.76	6.80	6.95	7.02	7.08	7.11	7.12	7.11	7.15	7.26
	90 th	6.27	6.28	6.29	6.29	6.30	6.33	6.37	6.44	6.53	6.60	6.70	6.73	6.75	6.81	6.94
	75 th	5.83	5.84	5.85	5.85	5.86	5.89	5.94	6.01	6.10	6.18	6.30	6.34	6.38	6.47	6.61
	50 th	5.38	5.40	5.41	5.41	5.43	5.46	5.51	5.58	5.68	5.76	5.84	5.96	6.02	6.13	6.29
	25 th	4.94	4.96	4.97	4.97	4.99	5.02	5.07	5.15	5.25	5.34	5.50	5.57	5.65	5.78	5.97
	10 th	4.50	4.52	4.53	4.53	4.55	4.59	4.64	4.72	4.82	4.92	5.01	5.18	5.29	5.44	5.64
	3 rd	4.06	4.08	4.09	4.09	4.11	4.15	4.21	4.40	4.40	4.50	4.70	4.80	4.92	5.10	5.32
Height (cm)	102.2	107.1	114.5	123.0	128.6	133.9	138.6	144.0	150.8	156.8	160.8	163.7	164.8	164.9	165.9	166.3
Weight (kg)	15.5	17.1	18.8	22.2	25.2	28.1	31.9	34.4	39.5	44.6	49.5	52.3	54.8	55.8	56.2	57.5

PWV_{ao}, aortic pulse wave velocity.




FIGURE 2 Smoothed median curve (grey curve) and first derivative of smoothed median curve of aortic pulse wave velocity (PWV_{ao} ; black curve) related to age for boys (a) and girls (b).

potential change of the position of the aortic bifurcation during growth is concerned, on the basis of our literature search we could not find any data about the change of the position of the aortic bifurcation related to its vertebral position during growing between ages of 3 and 18 years. In the available anatomical textbooks the length and the position of aortic parts (thoracic, abdominal) are determined in the relationship with vertebral number (thoracic or abdominal). Minor changes in the position of the aortic bifurcation, related to its vertebral position during growing cannot be excluded theoretically. Furthermore, this kind of potential bias due to the growing may exist if the measurement of the aortic PWV would be performed by carotid-femoral method.

The overall increase with age of mean PWV_{ao} values was identical in both sexes, only biologically insignificant differences were observed between them. These findings suggest that no significant structural difference exists between sexes in this age range.

Regarding the augmentation index, Ayer *et al.* measured carotid augmentation index (AIx) in 8-year-old children to examine the influences of sex, height and arterial stiffness on central arterial pulse wave augmentation [34]. They found that carotid AIx was significantly higher in girls than boys and it was independent of height, carotid artery diameter and stiffness. In contrast in our study we focused on the PWV and proved that the PWV_{ao} does not increase




FIGURE 3 Standardized median curves of aortic pulse wave velocity (PWV_{ao}), mean arterial pressure (MAP) and brachial SBP related to age for boys (a) and girls (b).

constantly with age. In the very young age groups (between 3 and 8 years), the median PWV_{ao} values are practically constant for both sexes.

The second original finding was that we observed a very important (previously not described) characteristic of the smoothed percentile curves of PWV_{ao} values.

During a detailed and careful analysis when the first derivatives of the smoothed median curve of PWV_{ao} were applied, it emerged that the steepest increase differs in terms of time between boys and girls, that is it occurs at 12.1 years in boys and at 10.4 years in girls. We can hypothesize that this newly observed phenomenon might be explained by the differing onset of adolescence in boys and girls.

When we compared the age-related changes of the median of the standardized values of PWV_{ao} , MAP and SBP_{brach} , a further new and interesting feature appeared. We observed marked differences between the ages of 3 and 8 years in the trends of PWV_{ao} , MAP and SBP_{brach} . Interestingly, the PWV_{ao} remained practically unchanged, whereas the blood pressure trend constantly increased. This raises several physiological questions. We might suppose that,

at a very young age, the aortic wall is so elastic that an increase in blood pressure does not cause any consequential stiffening in the aortic wall, but, clearly, further, specifically designed studies would be needed to confirm this hypothesis. Similarly the practically identical increase of the standardized PWV_{ao} and peripheral blood pressure at the later ages of 10–18 years might be due to a gradual loss of aortic elastic properties.

Several authors have provided evidence about the changes in the structure and composition of the aortic wall at an early age, specifically that ageing causes a loss in elastic properties [35–39]. It is possible that the continuously increasing blood pressure which occurs between the ages of 3 and 18 years may lead to a simultaneous increase in lateral tension in the aortic wall. Then beyond the age of 10–12 years, this load cannot be compensated for as successfully as was the case between 3 and 8 years. Hence, the rise in pressure will inevitably be accompanied by a rise in aortic stiffness.

Whether the abnormally steep increase of PWV_{ao} (e.g. an individual's increase in PWV_{ao} from the 25th to 90th percentile within a few years) might have pathological value in assessing cardiovascular risk characterized by the accelerated stiffening of the aortic wall is something which remains to be clarified by further, preferably longitudinal examination.

Assessing our age-related mean PWV_{ao} values in the light of earlier studies, the observed absolute values and their changes in relation to age seem to be very similar, suggesting that the PWV_{ao} values provided by the oscillometric method fall strictly within the range found by other examiners (Table 1). Unfortunately, a more detailed comparison of the observed mean PWV_{ao} values between studies cannot be undertaken, as neither the values of the measured distances nor the transit times of the pulse pressure waves between the observational points are given by the cited authors.

In this respect, our measured mean PWV_{ao} values (5.5–6.5 m/s in boys and 5.6–6.4 m/s in girls) are closest to the results of Reusz *et al.* published recently, although the upper age of their participants was 20 years [24]. Their applied distance measurement [(femoral site–jugular notch) – (carotid site–jugular notch)] and our distance [Jug-Sy, the distance between the sternal notch (jugulum) and upper part of the pubic bone (symphysis)] differs markedly. To be able to judge the difference in distance and to be able to compare the results more precisely, we carried out a small sub-study on our population involving 56 subjects aged between 6 and 18 (quite well distributed in terms of age). We measured the distances in the same subject by both methods and found a difference of 4.6 cm, i.e. the Jug-Sy distance was 4.6 cm longer on average than the [(femoral site–jugular notch) – (carotid site–jugular notch)] distance. Consequently, the PWV_{ao} differences between the data of Reusz *et al.* [24] and of ourselves (roughly 1 m/s) originated basically from the different methods which were used.

To summarize our study, we provided PWV_{ao} reference values for 3374 healthy whites, reasonably balanced in terms of age from 3 to 18 years, using a valid and reliable technique which can be used even at very young ages. We

first detailed the specific characteristics of the changes of PWV_{ao} – namely that the PWV_{ao} remains unchanged in the very young (despite increasing blood pressure), and we found the gradual increase of PWV_{ao} in adolescence – which is proportionate to the increase in blood pressure.

Study limitations

Our data are collected only in white population with no any ethnic or race mixture. Data collection in other population may be the issue of future researches.

ACKNOWLEDGEMENTS

The authors thank János Hubert, Györgyi Jákli, Ilona Kelemen, Hajnalka Klemencz (data collection) and Gábor ÉRSEK (data analysis) for their contribution to this work.

This study was supported by the Hungarian National Research Foundation (OTKA) No.78480.

Conflicts of interest

M.I. owns shares in TensioMed Ltd., the company which manufactures the Arteriograph and other devices for measuring vascular stiffness. F.T.M. is employed at TensioMed Ltd. The other authors have declared no conflicts of interest.

REFERENCES

- Mancia G, De Backer G, Dominiczak A, Cifkova R, Fagard R, Germano G, *et al.* 2007 ESH-ESC practice guidelines for the management of arterial hypertension: ESH-ESC Task Force on the Management of Arterial Hypertension. *J Hypertens* 2007; 25:1751–1762.
- Celermajer DS, Sorensen KE, Gooch VM, Spiegelhalter DJ, Miller OI, Sullivan ID, *et al.* Noninvasive detection of endothelial dysfunction in children and adults at risk of atherosclerosis. *Lancet* 1992; 340:1111–1115.
- Meyer AA, Kundt G, Steiner M, Schuff-Werner P, Kienast W. Impaired flow-mediated vasodilation, carotid artery intima-media thickening, and elevated endothelial plasma markers in obese children: the impact of cardiovascular risk factors. *Pediatrics* 2006; 117:1560–1567.
- McEniery CM, Bolton CE, Fawke J, Hennessy E, Stocks J, Wilkinson IB, *et al.* Cardiovascular consequences of extreme prematurity: the EPICure study. *J Hypertens* 2011; 29:1367–1373.
- Wiegman A, de Groot E, Hutten BA, Rodenburg J, Gort J, Bakker HD, *et al.* Arterial intima-media thickness in children heterozygous for familial hypercholesterolaemia. *Lancet* 2004; 363:369–370.
- Riggio S, Mandraffino G, Sardo MA, Iudicello R, Camarda N, Imbalzano E, *et al.* Pulse wave velocity and augmentation index, but not intima-media thickness, are early indicators of vascular damage in hypercholesterolemic children. *Eur J Clin Invest* 2010; 40:250–257.
- Singh TP, Groehn H, Kazmers A. Vascular function and carotid intima-media thickness in children with insulin-dependent diabetes mellitus. *J Am Coll Cardiol* 2003; 41:661–665.
- Jarvisalo MJ, Raitakari M, Toikka JO, Putto-Laurila A, Rontu R, Laine S, *et al.* Endothelial dysfunction and increased arterial intima-media thickness in children with type 1 diabetes. *Circulation* 2004; 109:1750–1755.
- Litwin M, Trelewicz J, Wawer Z, Antoniewicz J, Wierzbicka A, Rajszyz P, *et al.* Intima-media thickness and arterial elasticity in hypertensive children: controlled study. *Pediatr Nephrol* 2004; 19:767–774.
- de Divitiis M, Pilla C, Kattenhorn M, Donald A, Zadinello M, Wallace S, *et al.* Ambulatory blood pressure, left ventricular mass, and conduit artery function late after successful repair of coarctation of the aorta. *J Am Coll Cardiol* 2003; 41:2259–2265.
- Ou P, Celermajer DS, Raisky O, Jolivet O, Buyens F, Herment A, *et al.* Angular (Gothic) aortic arch leads to enhanced systolic wave reflection, central aortic stiffness, and increased left ventricular mass late after aortic coarctation repair: evaluation with magnetic resonance flow mapping. *J Thorac Cardiovasc Surg* 2008; 135:62–68.

12. Sheth RD, Perez MD, Goldstein SL. Cardiovascular calcifications in paediatric patients receiving maintenance dialysis. *Pediatr Nephrol* 2003; 18:810–813.
13. Charakida M, Donald AE, Green H, Storry C, Clapson M, Caslake M, *et al*. Early structural and functional changes of the vasculature in HIV-infected children: impact of disease and antiretroviral therapy. *Circulation* 2005; 112:103–109.
14. Noto N, Okada T, Yamasuge M, Taniguchi K, Karasawa K, Ayusawa M, *et al*. Noninvasive assessment of the early progression of atherosclerosis in adolescents with Kawasaki disease and coronary artery lesions. *Pediatrics* 2001; 107:1095–1099.
15. Tedesco MA, Di Salvo G, Ratti G, Natale F, Calabrese E, Grassia C, *et al*. Arterial distensibility and ambulatory blood pressure monitoring in young patients with neuro-fibromatosis type 1. *Am J Hyperten* 2001; 14:559–566.
16. Cheung YF, Brogan PA, Pilla CB, Dillon MJ, Redington AN. Arterial distensibility in children and teenagers: normal evolution and the effect of childhood vasculitis. *Arch Dis Child* 2002; 87:348–351.
17. Tauzin L, Rossi P, Giusano B, Gaudart J, Boussuges A, Fraisse A, *et al*. Characteristics of arterial stiffness in very low birth weight premature infants. *Pediatr Res* 2006; 60:592–596.
18. Kwok KL, Ng DK, Cheung YF. BP and arterial distensibility in children with primary snoring. *Chest* 2003; 123:1561–1566.
19. Tedesco MA, Ratti G, Di Salvo G, Martiniello AR, Limongelli G, Grieco M, *et al*. Noninvasive evaluation of arterial abnormalities in young patients with neuro-fibromatosis type 1. *Angiology* 2000; 51:733–741.
20. Ahimastos AA, Formosa M, Dart AM, Kingwell BA. Gender differences in large artery stiffness pre and postpuberty. *J Clin Endocrinol Metab* 2003; 88:5375–5380.
21. Avolio AP, Deng FQ, Li WQ, Luo YF, Huang ZD, Xing LF, *et al*. Effects of aging on arterial distensibility in populations with high and low prevalence of hypertension: comparison between urban and rural communities in China. *Circulation* 1985; 71:202–210.
22. Avolio AP, Chen SG, Wang RP, Zhang CL, Li MF, O'Rourke MF. Effects of aging on changing arterial compliance and left ventricular load in a northern Chinese urban community. *Circulation* 1983; 68:50–58.
23. Kis E, Cseperekal O, Horvath Z, Katona G, Fekete BC, Hrapka E, *et al*. Pulse wave velocity in end-stage renal disease: influence of age and body dimensions. *Pediatr Res* 2008; 63:95–98.
24. Reusz GS, Cseperekal O, Temmar M, Kis E, Cherif AB, Thaleb A, *et al*. Reference values of pulse wave velocity in healthy children and teenagers. *Hypertension* 2010; 56:217–224.
25. Urbina E, Williams R, Alpert BS, Collins RT, Daniels SR, Hayman L, *et al*, on behalf of the American Heart Association Atherosclerosis, Hypertension, and Obesity in Youth Committee of the Council on Cardiovascular Disease in the Young Noninvasive Assessment of Subclinical Atherosclerosis in Children and Adolescents. Recommendations for standard assessment for clinical research: a scientific statement from the American Heart Association. *Hypertension* 2009; 54:919–951.
26. Rosario AS, Kurth BM, Stolzenberg H, Ellert U, Neuhauser H. Body mass index percentiles for children and adolescents in Germany based on a nationally representative sample (KiGGS 2003–2006). *Eur J Clin Nutr* 2010; 64:341–349.
27. National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics* 2004; 114 (2 Suppl 4): 555–576.
28. Baulmann J, Schillings U, Rickert S, Uen S, Dusing R, Illyes M, *et al*. A new oscillometric method for assessment of arterial stiffness: comparison with tonometric and piezo-electronic methods. *J Hypertens* 2008; 26:523–528.
29. Horvath IG, Nemeth A, Lenkey Z, Alessandri N, Tufano F, Kis P, *et al*. Invasive validation of a new oscillometric device (Arteriograph) for measuring augmentation index, central blood pressure and aortic pulse wave velocity. *J Hypertens* 2010; 28:2068–2075.
30. Sugawara J, Hayashi K, Yokoi T, Tanaka H. Age-associated elongation of the ascending aorta in adults. *J Am Coll Cardiol Cardiovasc Imag* 2008; 1:739–748.
31. Mueller WH, Martorell R. Reliability and accuracy of measurement. In: Lohman TG, Roche AF, Martorell R, editors. *Anthropometric standardization reference manual*. Champaign, Illinois: Human Kinetics Books; 1988. pp. 83–86.
32. Pan H, Cole TJ. LMS Chartmaker, a programme to construct growth references using the LMS method. 2010. Version 2.43. <http://www.healthforallchildren.co.uk/>.
33. Cole TJ, Green PJ. Smoothing reference centile curves: the LMS method and penalised likelihood. *Stat Med* 1992; 11:1305–1319.
34. Ayer JG, Harmer JA, Marks GB, Avolio A, Celermajer DS. Central arterial pulse wave augmentation is greater in girls than boys, independent of height. *J Hypertens* 2010; 28:306–313.
35. Martyn CN, Greenwald SE. Impaired synthesis of elastin in walls of aorta and large conduit arteries during early development as an initiating event in pathogenesis of systemic hypertension. *Lancet* 1997; 350:953–955.
36. Glukhova MA, Frid MG, Koteliansky VE. Phenotypic changes of human aortic smooth muscle cells during development and in the adult vessel. *Am J Physiol* 1991; 261 (4 Suppl):78–80.
37. Van Meurs-Van Woezik H, Klein HW, Markus-Silvis L, Krediet P. Comparison of the growth of the tunica media of the ascending aorta, aorta isthmus and descending aorta in infants and children. *J Anat* 1983; 136:273–281.
38. Senzaki H, Akagi M, Hishi T, Ishizawa A, Yanagisawa M, Masutani S, *et al*. Age-associated changes in arterial elastic properties in children. *Eur J Pediatr* 2002; 161:547–551.
39. Niwa K, Perloff JK, Bhuta SM, Laks H, Drinkwater DC, Child JS, *et al*. Structural abnormalities of great arterial walls in congenital heart disease: light and electron microscopic analyses. *Circulation* 2001; 103:393–400.

Reviewers' Summary Evaluations

Reviewer 1

Aortic pulse wave velocity (aPWV) has been determined in one of the largest investigations undertaken in children of ages 3 to 18 years. aPWV is constant in the first decade (average of 5.55 m/s) but increases by 16% by age 18. However, this occurs some 2 years earlier in girls (10.4 years) compared with boys (12.1 years). This is consistent with studies showing girls having a higher carotid augmentation

index compared with boys at the same age of 8 years. One of the caveats of the study is that aPWV was not measured, but estimated from brachial cuff suprasystolic waves.

Reviewer 2

This article provides reference ranges for PWV in very young children. The data are of interest and adds to the literature on the progression of PWV with age but with the novelty of a detailed study in very young children.